

ACUTE TRANSFUSION REACTIONS

Recognise. Respond. Report.

PATIENT HAS SIGNS AND SYMPTOMS SUGGESTIVE OF POTENTIAL TRANSFUSION REACTION



Assess: rapid clinical assessment

Check: confirm patient ID band matches blood swing label details

Inspect: visual check of unit for turbidity, clots or abnormal appearance

Talk with the Patient: establish status, inform and comfort

Are symptoms **LIFE THREATENING?** Airway/Breathing/Circulation?
OR Wrong Blood Given? OR Evidence of Abnormal Unit?

YES

NO

Severe or Life Threatening Events

- ✓ **CALL** for **urgent** medical help and review
- ✓ **INITIATE** Resuscitation: ABC
- ✓ **DISCONNECT** IV infusion set/unit – do **NOT** discard/restart
- ✓ **MAINTAIN** venous access with saline via **NEW** infusion set
- ✓ **ADMINISTER** IV fluids/O₂ if clinically indicated
- ✓ **MONITOR** TPR/BP/SpO₂/urine output (q5-15 min)
- ✓ **TREAT** according to clinical status/symptoms, noting:
 - ? **anaphylaxis/severe allergy**: use NZRC Anaphylaxis Guide
 - ? **septic shock**: use DHB Sepsis Guidelines
 - ? **acute haemolysis**: maintain BP, force diuresis, alkalinise urine
 - ? **circulatory overload**: diuretics, O₂, positive airway pressure
 - ? **TRALI**: respiratory support, ask NZBS to start donor review

ALERT: Is haemorrhage a possible cause of the hypotension?
Resuscitate with fluids and consider further transfusion

INFORM your local clinical haematologist or TMS via Blood Bank ASAP or, contact directly if treatment advice needed

INFORM medical staff – seek **PROMPT** clinical review

Moderate Events

- ✓ All symptoms that are not classified as mild, severe or life threatening

Management

- ✓ Disconnect IV infusion set/unit – do **NOT** discard set/unit
- ✓ Replace IV infusion set
- ✓ Maintain venous access with saline
- ✓ Treat according to clinical status
- ✓ Do **NOT** restart transfusion

Mild Events

- ✓ Fever > 38°C and < 1.5°C above baseline with no other symptoms
- ✓ Localised rash with no other symptoms

Medical Review

- ? If fever – consider antipyrexial
- ? If localised rash – consider antihistamine

Management

- ✓ Consider restarting transfusion at slower rate. Directly observe for first 15 minutes
- ✓ Increase frequency of monitoring vital signs (TPR/BP/SpO₂) thereafter

Reporting

- ✓ Document in clinical notes
- ✓ Send NZBS ATR Notification Form (111F009) to Blood Bank
- ✓ No blood tests required

Investigations and Reporting

- ✓ **DO** – ‘Standard ATR Investigations’ and undertake ‘Additional Investigations’ as needed (*see below*)
- ✓ **COMPLETE** – NZBS ATR Notification Form (111F009)
- ✓ **SEND** – blood unit/IV set, ATR Notification Form and EDTA (pink) sample to BB and other samples to Pathology
- ✓ **RECORD** – in clinical notes

If symptoms worsen?

STOP transfusion and manage as per a **moderate** or **severe** event

ACUTE TRANSFUSION REACTIONS (ATR) – INFORMATION FOR CLINICAL STAFF

1. Recognise.

Signs and symptoms may include:

- Fever, chills, rigors
- Tachycardia, arrhythmias
- Hyper or hypotension, collapse
- Generalised flushing
- Rash, urticaria, angioedema
- Anxiety, severe apprehension
- Nausea, vomiting
- Pain (*chest, loin, muscle, bone, abdominal, cannula site/vein*)
- Dyspnoea, respiratory distress, hypoxia
- Pink/red/black urine or abnormal bleeding

2. Respond.

Management and investigations

- 1 **Clinical Review/Treatment** – as above
- 2 **Standard ATR Investigations** – for **ALL moderate and severe events** are:
 - EDTA (pink top) for serology – to Blood Bank
 - Full blood count/film and UE – to Pathology
 - Ward urinalysis for blood/haemoglobin
- 3 **Additional Investigations:** if...
 - ? **Haemolysis**: consider haptoglobin, LDH, coag’s
 - ? **Respiratory Distress**: consider CXR, ABGs, BNP
 - ? **Sepsis/Shock**: consider blood cultures from patient
 - ? **Severe allergy/anaphylaxis**: consider serum tryptase & query need for anti-IgA antibodies

3. Report.

Haemovigilance to Blood Bank (BB)

- Report all mild, moderate and severe ATRs using the NZBS ATR Notification Form
- If the event is **moderate** or **severe**, remember to include the EDTA (pink) sample, the discontinued unit and blood IV infusion set with the ATR Form (111F009)

Advice on management, further transfusion needs or recurrent reactions should be discussed with the transfusion medicine specialist (TMS) or the clinical haematology consultant.